

Incidence of Delayed Intracranial Hemorrhage in Patients Taking Warfarin that Sustain Head Trauma

Lauren Segal DO, Brian Collins DO, Maricel Dela Cruz DO, Michael Baier DO, Scott Plasner DO

Aria Jefferson Health, Department of Emergency Medicine, Philadelphia, PA, United States

INTRODUCTION

With an aging United States population, individuals taking anticoagulant medications, specifically warfarin, are common. A large proportion of the elderly and those in lower socioeconomic classes utilize warfarin due to its affordability. Emergency medicine physicians are challenged in treating these patients after they sustain head trauma. It is agreed that the overall mortality in anticoagulated patients is significantly higher in patients on warfarin versus non-anticoagulated patients (6, 12). Of particular interest is the incidence of delayed intracranial hemorrhage in patients with blunt head trauma while on warfarin. To date, there is limited data on this matter.

MATERIALS AND METHODS

A retrospective observational study was conducted using the following software: Wellsoft ED charts, Allscripts Sunrise inpatient data and radiographic studies on PACS from January 2009 to January 2012 at a Level II community trauma center, Aria Health Torresdale in Philadelphia, Pennsylvania. The Aria Health IRB approved this study. Participants included individuals admitted to the trauma service that were taking warfarin, suffered blunt head trauma, and had an initially negative CT Brain. Patients had to be 18 years or older to be eligible. There were no interventions performed, as this was a retrospective analysis.

CONCLUSIONS

There is a risk of delayed intracranial hemorrhage in individuals using warfarin. This study found that none of the three patients with delayed intracranial hemorrhage had a significant change in their clinical status. No neurosurgical intervention was required. While there is a slight risk of delayed hemorrhage in this patient population, the most appropriate management may be to monitor these patients for clinical status change and only then could the decision for repeat imaging be made. On a systems-wide scale, this has the potential to decrease spending on imaging and reduce unnecessary radiation exposure.





Figure
Normal Computed Tomography of the Brain.



REFERENCES

Franko, J., Kish, K., O'Connell, B., Subramanian, S., Yuschak, J. Advanced age and preinjury warfarin anticoagulation increase the risk of mortality after head trauma. *The Journal of Trauma Injury, Infection, and Critical Care*. [Internet] 2006. [Cited 2014 April 19]. Available from: doi:10.1097/01.ta.0000224220.89528.fc.

McMillian, W.D., Rogers, F.B. Management of prehospital antiplatelet and anticoagulant therapy in traumatic head injury: a review. *The Journal of Trauma Injury, Infection, and Critical Care*. [Internet] 2009. [Cited 2014 April 19]. Available from: doi:10.1097/TA.0b013e3181978e7b.

Brewer, E.S., Reznikov, B., Liberman, R.F., Baker, R.A., Rosenblatt, M.S., David, C.A., Flacke, S. Incidence and predictors of intracranial hemorrhage after minor head trauma in patients taking anticoagulant and antiplatelet medication. *The Journal of Trauma Injury, Infection, and Critical Care*. [Internet] 2011. [Cited 2014 April 19]. Available from: doi:10.1097/TA.0b013e3181e5e286.

Claudia, C., Claudia, R., Agostino, O., Simone, M., Stefano, G. Minor head injury in warfarinized patients: indicators of risk for intracranial hemorrhage. *The Journal of Trauma Injury, Infection, and Critical Care*. [Internet] 2011. [Cited 2014 April 19]. Available from: doi:10.1097/TA.0b013e318203lab7.

Cohen, D.B., Rinker, C., Wilberger, J.E. Traumatic brain injury in anticoagulated patients. *The Journal of Trauma Injury, Infection, and Critical Care*. [Internet] 2006. [Cited 2014 April 19]. Available from: doi:10.1097/01.ta.0000196542.54344.05.

Hill, J. Delayed intracranial hemorrhage in the anticoagulated blunt trauma patient: routine repeat head computed tomography is unnecessary. *2013. Critical Care Medicine*, 41(12).

Howard, J.L., Cipolle, M.D., Horvat, S.A., Sabella, V.M., Reed, J.F., Fulda, G., Tinkoff, G., Pasquale, M.D. Preinjury warfarin worsens outcome in elderly patients who fall from standing. [Internet] 2009. [Cited 2014 April 19]. Available from: doi:10.1097/TA.0b013e3181a59728.

Ivascu, F.A., Janczyk, R.J., Junn, F.S., Bair, H.A., Bendick, P.J., Howells, G.A. Treatment of trauma patients with intracranial hemorrhage on preinjury warfarin. *The Journal of Trauma Injury, Infection, and Critical Care*. [Internet] 2006. [Cited 2014 April 19]. Available from: doi:10.1097/01.ta.0000223944.25922.91.

Joseph, B., Sadoun, M., Aziz, H., Tang, A., Wynne, J.L., Pandit, V., Kulvatunyou, N., O'Keefe, T., Friese, R.S., Rhee, P. Repeat head computed tomography in anticoagulated traumatic brain injury patients: still warranted. *The American Surgeon*. [Internet] 2014. [Cited 2014 April 19]. Available from: EBSCO Publishing IP: 202.177.173.189.

Nishijima, D.K., Offerman, S.R., Ballard, D.W., Vinson, D.R., Chettipally, U.K., Rauchwerger, A.S., Reed, M.E., Holmes, J.F. Immediate and delayed traumatic intracranial hemorrhage in patients with head trauma and pre-injury warfarin or Clopidogrel use. *Annals of Emergency Medicine*. [Internet] 2012. [Cited 2014 April 19]. Available from: doi:10.1016/j.annemergmed.2012.04.007.

OBJECTIVES

The objective of this study was to determine the incidence of delayed intracranial hemorrhage by computed tomography in individuals taking warfarin. It is hypothesized that the incidence of delayed intracranial hemorrhage in this specific population is low. As a result, this data can potentially prevent unnecessary admissions to the hospital, decrease radiation exposure, and reduce costs.

RESULTS

The primary outcome of this study was to determine the incidence of delayed intracranial hemorrhage in patients sustaining blunt head trauma while anticoagulated on warfarin from 2009 to 2012. In this chart review study, there were several thousand patients admitted to the trauma service during the time span of interest. Of those, 73 patients were admitted to the trauma service that satisfied this study's inclusion criteria. There were a multitude of blunt traumatic mechanisms of injury, including but not limited to, accidental falls, syncope, assaults, and MVCs. These patients were anticoagulated on warfarin for a variety of reasons, most commonly for atrial fibrillation. In this cohort, there were three patients who had an initially negative CT Brain that then developed an intracranial hemorrhage as demonstrated upon a 6-hour follow-up CT scan. The remaining 70 patients had negative imaging on a 6-hour follow-up CT Brain.

The authors of this presentation do not have any disclosures.